

To appear in The Astrophysical Journal Letters

Surface Brightness and Stellar Populations at the Outer Edge of the Large Magellanic Cloud: No Stellar Halo Yet

Carme Gallart¹

Instituto de Astrofísica de Canarias. 38200 La Laguna. Tenerife, Canary Islands. Spain.

`carme@iac.es`

Peter B. Stetson

Dominion Astrophysical Observatory, Herzberg Institute of Astrophysics, National Research Council of Canada, 5071 West Saanich Road, Victoria, BC V9E 2E7, Canada

`Peter.Stetson@nrc-cnrc.gc.ca`

Eduardo Hardy^{2,3}

National Radio Astronomy Observatory. Casilla 36-D, Santiago, Chile

`ehardy@nrao.edu`

Frederic Pont

Observatoire de Genève, CH-1290 Sauverny, Switzerland

`Frederic.Pont@obs.unige.ch`

and

Robert Zinn

Department of Astronomy, Yale University, P.O. Box 208101, New Haven, CT 06520-8101

`zinn@astro.yale.edu`

¹Ramón y Cajal Fellow.

²The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc.

³Adjoint Professor, Astronomy Dept. Universidad de Chile

ABSTRACT

We present a high quality CMD for a $36' \times 36'$ field located 8° ($\simeq 7$ kpc) from the LMC center, as well as a precise determination of the LMC surface brightness derived from the resolved stellar population out to this large galactocentric radius. This deep CMD shows for the first time the detailed age distribution at this position, where the surface brightness is $V \simeq 26.5$ mag/ $\square''$. At a radius $R \simeq 474'$ the main sequence is well populated from the oldest turnoff at $I \simeq 21.5$ to the 2.5 Gyr turnoff at $I \simeq 19.5$. Beyond this radius, a relatively strong gradient in the density of stars with ages in the $\simeq 2.5$ –4 Gyr range is apparent. There are some stars brighter and bluer than the main population, quite uniformly distributed over the whole area surveyed, which are well matched by a 1.5 Gyr isochrone and may be indicative of a relatively recent star formation, or merger, event. The surface brightness profile of the LMC remains exponential to this large galactocentric radius and shows no evidence of disk truncation. Combining the information on surface brightness and stellar population we conclude that the LMC disk extends (and dominates over a possible stellar halo) out to a distance of at least 7 kpc. These results confirm that the absence of blue stars in the relatively shallow off-center CMDs of dIrr galaxies is not necessarily evidence for an exclusively old stellar population resembling the halo of the Milky Way.

Subject headings: galaxies: Magellanic Clouds—galaxies: halos—galaxies: evolution—galaxies: stellar content

1. Introduction

The nature of the stellar populations at the outer edges of dwarf irregular (dIrr) galaxies, and in particular whether these galaxies have a truly old stellar halo has been a subject of some controversy over the last few years. The presence of a background sheet of red stars, extending beyond the most prominent blue, young stellar population in IC1613 and other dIrr galaxies was discussed already by Baade (Sandage 1971). But, what is the nature of this sheet of red stars? Minniti & Zijlstra (1996) claimed that the Local Group dIrr WLM has an old (as old as the Milky Way globular clusters) stellar halo, from the sole evidence that the color-magnitude diagram (CMD) of an outer field extending out to $8'$ from the center of the galaxy, along its minor axis, does not contain blue stars down to $I \simeq 23$ ($M_I \simeq -1.8$), and shows a well defined tip of the RGB. However, since all stars older than about 1 Gyr produce a RGB at very approximately the same location, and main-sequence (MS) stars

older than a few hundred Myr would be below the limit of Minniti & Zijlstra’s photometry, all that can be confidently concluded from the lack of bright MS stars is that the stellar population is older than a few hundred Myr (see also the discussion by Aparicio, Tikhonov & Karachentsev 2000). Other evidence, such as the presence of C stars in the outer regions of dwarf galaxies (e.g. Letarte et al. 2002), or a full modeling of the CMD using synthetic CMDs (Aparicio & Tikhonov 2000) suggests in fact the presence of a range of ages.

Determining the age structure and kinematic properties of the outer red component of dIrr galaxies may hold important clues on the process of dwarf galaxy formation and evolution. Do these stars represent a stellar component truly differentiated from the younger one, with the characteristics of a halo population similar to that of the Milky Way (old, metal poor, devoid of gas, kinematically hot), or is there a smooth gradient of stellar ages, which might suggest that star formation has been “shrinking” with time? (Hidalgo, Marín-Franch & Aparicio 2003)

A complete answer to these questions requires both very deep photometry, reaching the oldest MS turnoffs, and relatively high dispersion spectroscopy to obtain kinematics of individual stars. From deep CMDs at different galactocentric distances it is possible to characterize the ages of the stellar populations present, and to assess, therefore, whether a “break” in the stellar population, indicating an edge to the zone of protracted star formation and a transition to an old stellar halo actually occurs. The observational requirements for this task are, however, challenging. For the Milky Way satellites, it is necessary to reach $V \simeq 24$ to obtain good precision photometry of the old MS turnoff region. For the rest of the Local Group, $V \simeq 28$ is necessary. While the first is attainable from the ground, the latter is only possible using the HST equipped with ACS. Obtaining the required spectroscopic information is a highly challenging enterprise.

The Magellanic Clouds are the two nearest Irr galaxies. Their old MS turnoffs are located near $V \simeq 23$. While this magnitude is easily attainable from the ground, the huge angular sizes of the Clouds makes the mapping of their full extent and the characterization of their stellar population gradients a daunting task. As discussed by Irwin (1991) on the basis of isopleth maps, the LMC extends over an elliptical area covering 23° by 17° , with the major axis oriented roughly in the North-South direction. This is also basically the extent of the LMC cluster system. On the North side, North of declination -62° ($\simeq 8^\circ$ from the center), the isopleths abruptly separate from each other. Does this indicate some kind of break in the characteristics of the stellar populations? Some information about the ages of the stars present at this position comes from the study of Kunkel, Irwin & Demers (1997), who searched for C stars in the periphery of the two Magellanic Clouds. Interestingly, they found a relatively sharp boundary for the outermost distribution of C stars at a radius of $\simeq 8^\circ$

along the major axis, which may indicate the actual maximum extent of an intermediate-age population. A detailed characterization of such a population, however, requires obtaining CMDs reaching the oldest MS turnoffs. A full dynamical study of the C star population of the LMC based on 1043 objects out to 8° from the center was published by van der Marel et al. (2002), who inferred the presence of a dark-matter halo containing roughly 2/3 of the total mass of the LMC. The presence of a kinematically hot, metal-poor, old stellar halo has been inferred by Minniti et al. (2003) from the kinematics of a sample RR Lyrae stars.

With the purpose of characterizing the stellar population gradients in the LMC, we imaged four fields situated at distances ranging from 3.1 to 8 degrees from the LMC center using the CTIO 4m MOSAIC wide field imager. The full analysis of the whole dataset is underway (Gallart et al. 2004). In this Letter we analyze the CMD of the outermost field, which shows a somewhat surprising result on the age composition of the outer LMC, and use the whole dataset to trace the surface brightness across this galaxy.

2. Observations and data reduction.

We obtained V and I images of a LMC field centered at 7.8° from its center to the North ($\alpha : 05:13:17$; $\delta : -61:58:40$) on January 17th and 18th, 2001, using the Mosaic II CCD Imager on the CTIO Blanco 4m telescope. The Mosaic II provides a total field of $36' \times 36'$, with a resolution of $0''.27/\text{pixel}$. Seeing was typically around $1''.0$. Exposure times were 6×900 s in V *plus* 8×600 s in I. Several short exposures were also obtained to retrieve the photometry of the brightest stars. The nights were photometric, and the Landolt (1992) standard star fields SA92, SA95, SA98, SA104 were observed several times each night for calibration purposes. We used Stetson's (2000) photometry for these fields to calibrate our photometry. During the same run and in another one in December 1999, we obtained photometry of similar quality for three more internal fields. These results will be described elsewhere (Gallart et al. 2004), and used occasionally here (Section 3).

The Mosaic frames were reduced in a standard way, as indicated in the *CTIO Mosaic II Imager User Manual* (Schommer et al. 2000), using the MSCRED package within IRAF. Finally, profile-fitting photometry of the LMC fields was obtained using the DAOPHOT / ALLFRAME suite of computer programs (Stetson 1987, 1994). A full description of the data reduction and photometry is provided in Gallart et al. (2004).

3. The LMC surface brightness profile: reaching the outer edge?

We have derived the surface brightness for the four fields we observed in the LMC, by integrating the flux in the CMD¹. This method has the advantage that a very accurate field subtraction can be performed at the very low surface-brightness regime as foreground stars and background galaxies are for the most part well separated from the characteristic sequences of the LMC CMD. This is particularly important in the outermost field, where the flux contribution of Milky Way foreground objects exceeds by far that of the galaxy. The process will be fully described in Gallart et al. (2004). In short, we used the predictions of the Besançon Galaxy model (Robin et al. 2003), but scaling the number of predicted Milky Way stars to the number observed in our field in zones in the CMD clearly devoid of LMC stars. Without this scaling, we found that Besançon models predicted 25% more Galactic stars than we observed at relatively bright magnitudes. In addition, we only considered stars fainter than $V = 15$ when integrating the flux in the field, since no stars belonging to the LMC were observed brighter than this limit. Since field stars brighter than $V = 15$ are the main contributor to the total flux of the field, this greatly reduced the error in its subtraction. We neglected the contribution of background unresolved galaxies, since they are faint.

In Figure 1, the surface brightness of our fields measured in V (dots) is represented as a function of distance from the center of mass of the LMC. We have divided the outermost field into seven sections at increasing distances from the center of the LMC and calculated the surface brightness for each, in order to check whether the surface brightness variation remained smooth (which seems to be the case) or whether some kind of truncation was observed instead. The error bars were estimated by repeating the calculations with sky backgrounds that varied by 25% from our best estimate. We have added (triangles) two measurements from Hardy (1978). Exponential laws for the LMC disk as calculated from the whole dataset (Hardy *plus* our data, solid line), and as derived by Hardy (long-dashed line) and Bothun & Thompson (1988, short-dashed line) are represented. Our fit has a slope similar to that given by Bothun & Thompson, but is brighter. We also don't see evidence of the truncation found by these authors at a galactocentric distance of approximately $170'$. We conclude that the surface-brightness measurements displayed are consistent with an exponential law out to the last point measured.

¹The contribution to the total flux of the unresolved LMC population (i.e. by the stars fainter than $M_V = 5$) may range between 1 and 10% of the total flux, as estimated from synthetic CMDs for a composite population (constant star formation rate from 15 Gyr ago to the present time) and for a purely old population. We estimate, therefore, that we may be underestimating the surface brightness by a maximum of 0.1 magnitudes.

The field we will discuss in this Letter is the outermost one, situated from 450' to 490' from the center of the LMC. The surface brightness in this field ranges from 26.2 to 26.8 mag/□'' in V and is one magnitude brighter in I . To put the location of our field in context, we have drawn an arrow at the location of the outermost radius for which the LMC surface brightness was measured by Bothun & Thompson (1988). The tip of the arrow indicates an estimate of the V magnitude measured by these authors at this position, calculated using their B magnitude and assuming a color $(B - V) = 0.5$. This location also approximately coincides with the outer radius of the HI measured by Kim et al. (1998). The radius of the outermost field that we have measured is roughly equal to the radii of the maximum extension of the LMC cluster system (Olszewski et al. 1988) and the outer edge of the C-star system obtained by Kunkel et al. (1997).

4. The Color-Magnitude Diagrams

Figure 2 shows $[(V - I), I]$ CMDs of the LMC, in order of increasing distance, R , from the LMC center, from galactocentric radius 450 to 490' (see caption for details). Stars with at least one valid measurement in each filter have been selected using the following limits for the error and shape parameters given by ALLFRAME: $\sigma_{(V-I)} (= \sqrt{\sigma_V^2 + \sigma_I^2}) \leq 0.1$, $|sharp| \leq 0.3$ and $\chi \leq 4.5$. A total of 55000 stars down to $I \leq 24.0$ have been measured. The CMDs reach a couple of magnitudes below the old MS turnoff and therefore, this point is always cleanly reached with good photometric accuracy. Isochrones from Pietrinferni et al. (2004) for $Z=0.001$ and ages 13.5 and 8 Gyr, $Z=0.002$ and age 4.5 Gyr and $Z=0.004$ and ages 2.5 and 1.5 Gyr have been superimposed.

These CMDs show for the first time the details of the age structure of the stellar population at the outer edge of the LMC. Out to a radius $R \simeq 474'$ the MS appears populated quite smoothly from the oldest turnoff at $I \simeq 21.5$ to the 2.5 Gyr turnoff at $I \simeq 19.5$. Beyond this radius, a relatively strong gradient in the density of stars of ages $\simeq 4$ -2.5 Gyr begins to be apparent, with very few stars in this range beyond $R=480'$. There is a number of stars brighter and bluer than the main population, which are well matched by a 1.5 Gyr isochrone. They are distributed quite uniformly over the whole field. No foreground stars or background galaxies, however, are expected to populate the CMD at this position, and therefore they must be young stars genuinely belonging to the LMC. They may indicate a small burst of star formation 1.5 Gyr ago, the spatial extent of which cannot be evaluated with the current dataset. A survey of a larger area, and more fields at similar radius but different position angles would be necessary to ascertain whether it has been a global or a localized event. The fainter stars in the subgiant-branch region are well matched by the 13.5

Gyr, $Z=0.001$ isochrone, but no blue extended horizontal branch can be seen in the CMDs. The red clump—or red horizontal branch—presents, however, a short faint blue extension.

Figure 3 displays the color function of the CMD integrated over the magnitude range $21.5 \leq I \leq 17.5$, i.e. including all stars from just below the red clump to the location of the oldest MS turnoffs. The star counts have been normalized to the number of stars in the last two bins, which mainly include the red clump and a portion of the RGB just below it. Since these structures contain stars of ages greater than $\simeq 1$ Gyr, this is equivalent to a normalization by an indicator of the mean star formation rate. The gradient in the number of blue stars present in these CMDs, and their color distribution is very striking, notably more pronounced than would be expected from a visual inspection of the CMDs alone. The main population of age ≥ 2.5 Gyr present in the CMD appears at $(V - I) \simeq 0.2$, and the non-negligible number of stars between $(V - I) = 0 - 0.2$ indicate the presence of the population of age $\simeq 1.5$ Gyr mentioned above (error-bars have been omitted for clarity, but they are smaller than the differences between the curves). Stars bluer than $(V - I) \simeq 0.5$ are younger than $\simeq 8$ Gyr, and it is in this interval of color that the largest differences between fields are apparent. Beyond $R \simeq 470'$ the population of stars younger than 8 Gyr starts to abruptly decrease. The interval $(V - I) = 0.6 - 0.7$ is dominated by stars older than 8 Gyr, for which the relative numbers are very similar at all radii. Differences in the color function appear again in the interval $(V - I) = 0.8 - 0.9$, which mainly contains the subgiant branches of populations of all ages.

5. Discussion

We have presented a high-quality CMD, reaching the oldest main sequence turnoffs, of an LMC field situated at $R=450-490'$ from the center of mass of the LMC, toward the North (i.e., in the direction of its major axis). Very few CMDs of the LMC field have been published beyond this radius. The most targeted, detailed work is, to our knowledge, that of Stryker (1984) who did photographic photometry of stars in an area of $118'4 \times 118'4$ situated 9° northeast of the LMC bar. She just reached the oldest MS turnoffs. In spite of the very large photometric errors, she was able to show that a blue population brighter than the old turnoff was indeed present. In the very large area surveyed, she was also unable to find evidence of a blue horizontal branch.

In this paper, we show the presence of a substantial intermediate-age stellar population at $R = 8^\circ (\sim 7 \text{ kpc})$ from the center of the LMC. A strong gradient in this population, however, occurs in a range of just 0.5 kpc. The combination of an exponential surface brightness profile together with intermediate-age stars indicates the presence of a dominant

disk population even at these large galactocentric distances. It would be of great interest to investigate the stellar populations at even larger radial distances, to ascertain whether eventually a background old population appears and stays for a while (possibly indicating a true old *stellar* halo population that would need to be investigated further), or whether the actual outer edge of the LMC is smoothly reached with a continuing gradient of stellar populations. The lack of a well populated, blue extended horizontal branch, however, both in the field investigated here and in the larger, even more peripheral field studied by Stryker (1984) seems to indicate that the amount of *field* stellar population as old and metal poor as that of the Milky Way halo globular clusters and dSph galaxies is small in the LMC.

Turning now to the origin of the intermediate-age population in the outer fields, it is important to ask whether it could have formed at its current position, or it must have migrated from a more internal birth site (Hodge, Hunter & Oey 2003). Since the rotation of the galaxy may make it difficult for stars to move outwards, this population of stars may be a sign that the HI gas extended as far as this outer field as recently as a few Gyr ago and has since (gradually, thus the observed gradient), receded to the more central regions. Regarding the youngest $\simeq 1.5$ Gyr old population, which seems to be quite uniformly distributed over the field, an interaction between the LMC and the SMC, or the merger of the LMC with a smaller galaxy are attractive mechanisms for either in-situ formation, or accretion of stars in an apparently discrete age range.

Our results, in any case, are a demonstration that the lack of MS stars and the presence of a RGB with characteristics very similar to that of an old population cannot be taken as a proof of the existence of an exclusively old stellar population. The presence of a true stellar halo in dIrr galaxies remains therefore unproven, and it needs to be investigated with much deeper CMDs aided with spectroscopic data to characterize robustly the metallicities and the kinematics of the individual stars.

This research is part of a joint project between the Universidad de Chile and Yale University that is partially funded by the Fundación Andes. C.G. acknowledges partial support from the IAC (Project 3I1902), the Spanish Ministry of Science and Technology (Plan Nacional de Investigación Científica, Desarrollo e Investigación Tecnológica, AYA2002-01939) and from the European Structural Funds. R.Z. acknowledges the support of NSF grant AST-0098428.

REFERENCES

- Aparicio, A., Tikhonov, N., & Karachentsev, I. 2000, AJ, 119, 177
- Aparicio, A. & Tikhonov, N. 2000, AJ, 119, 2183
- Bothun, G.D. & Thompson, I.B. 1988, AJ, 96, 1988
- Gallart, C., Stetson, P.B., Pont, F., Zinn, R. & Hardy, E. 2004, in preparation
- Hardy, E. 1978, PASP, 90, 132
- Hidalgo, S., Marín-Franch, A. & Aparicio, A. 2003, AJ, 125, 1247
- Hodge, P.W., Hunter, D. & Oey S. 2003, PASP, 115, 273
- Irwin, M.J. 1991 in *The Magellanic Clouds*. IAU Symposium 148, 453 Ed. R. Haynes and D. Milne. Kluwer Academic Publishers, Dordrecht.
- Kim, S., Staveley-Smith, L., Dopita, M.A., Freeman, K.C., Sault, R.J., Kesteven, M.J., McConnell, D. 1998, ApJ, 503, 674
- Kunkel, W.E., Irwin M.J. & Demers, S. 1997, A&AS, 122, 463
- Landolt, A.U. 1992, AJ, 104, 340
- Letarte, B., Demers, S., Battinelli, P., Kunkel, W.E. 2002, AJ, 123, 832
- Minniti, D. & Zijlstra, A. 1996, ApJ, 467, L13
- Minniti, D., Borissova, J., Rejkuba, M., Alves, D.R., Cook, K.H. & Freeman, K.C. 2003, Sci, 301, 1508
- Olszewski, E.W., Harris, H.C., Schommer, R.A. & Catena, R.W. 1988, AJ, 95, 84
- Pietrinferni, A., Cassisi, S., Salaris, M. & Castelli, F. 2004, ApJ, in press
- Robin, A.C., Reylé, C., Derrière S. & Picaud S. 2003, A&A, 409, 523
- Sandage, A. 1971, ApJ, 166, 13
- Schommer, R., Smith, C., Olsen, K. & Walker, A. Eds. 2000: CTIO MOSAIC II Imager User Manual (<http://www.ctio.noao.edu/mosaic/manual/index.html>)
- Stetson, P.B. 1987, PASP, 99, 191

Stetson, P.B. 1994, PASP, 106, 250

Stetson, P.B. 2000, PASP, 112, 925

Stryker, L.L. 1984, ApJSS, 55, 127

van der Marel, R. P., Alves, D. R., Hardy, E., Suntzeff, N. B. 2002, AJ, 124, 2639




Fig. 1.— V (dots) surface brightness profile of the LMC as measured from our MOSAIC data. The field discussed in this paper is the outermost one. The surface brightness has been calculated across this field and given for seven sections at increasing distances from the center of the LMC. Two measurements of Hardy (1978, his innermost point and the outermost one of his “outer regions”) have been added as triangles. The outermost radius measured by Bothun & Thompson (1988) is marked with an arrow, which also indicates an estimate of the V surface brightness measured by them at this radius, assuming $(B - V) = 0.5$. Exponential laws for the LMC disk as calculated from the whole dataset represented in the figure (Hardy 1978 *plus* our data, solid line; $\mu_i(0)=21.16$, $\alpha=98'.7$) and as derived by Hardy (1978, long-dashed line; $\mu_i(0)=21.18$, $\alpha=86'.2$) and Bothun & Thompson (1988, short-dashed line; $\mu_i(0)=21.16$, $\alpha=100'.6$) are represented.




Fig. 2.— Observed $[(V-I), I]$ CMDs in order of increasing distance from the LMC center. The number of stars in each CMD is the same and, since the density of LMC stars decreases as a function of radius, δR varies for each CMD. The range in R of each CMD is labeled. Isochrones by Pietrinferni et al. (2004) for $Z=0.001$ and ages 13.5 and 8 Gyr (red), $Z=0.002$ and age 4.5 Gyr (green) and $Z=0.004$ and ages 2.5 and 1.5 Gyr (blue) have been superimposed.




Fig. 3.— Color function of the CMD integrated over the magnitude range $21.5 \leq I \leq 17.5$, for each of the CMDs shown in Figure 2. Solid, long-dashed, short-long dashed, short-dashed, dot-short dashed and dotted lines are in order of increasing galactocentric distance, respectively (see text for details).